

Taiwania*An International Journal of Biodiversity*, established 1947

Editors-in-Chief	Chang-Fu Hsieh, Wen-Yuan Kao
Associate Editors	Pei-Chun Liao, Ming-Jou Wu, Jenn-Che Wang, Roland Kirschner, En-Cheng Yang, Yiching Lin, Pei-Jen Lee Shaner
Managing Editor	Kuo-Hsiung Wang

Editorial Board

Grecebio Jonathan D. Alejandro ; Philippines	Serge Morand , France
David E. Boufford ; USA	Bai-Ling Lin ; Taiwan
Su-Hwa Chen ; Taiwan	Hiroyoshi Ohashi ; Japan
Peter L. Chesson ; USA	Arun K. Pandey ; India
Wen-Liang Chiou ; Taiwan	Chiou-Rong Sheue ; Taiwan
Chang-Hung Chou ; Taiwan	Shiuh-Feng Shiao ; Taiwan
Gabor Csorba ; Hungary	Kitichate Sridith ; Thailand
Sauren Das ; India	Hang Sun ; China
Christopher Roy Fraser-Jenkins ; Nepal	Iván A. Valdespino ; Panama
Tsung-Hsin Hsieh ; Taiwan	Chun-Neng Wang ; Taiwan
Jer-Ming Hu ; Taiwan	Ruijiang Wang ; China
Shih-Tong Jeng ; Taiwan	Jiunn-Tzong Wu ; Taiwan
Yeong-Choy Kam ; Taiwan	Wen-Jer Wu ; Taiwan
Chow-Yang Lee ; USA	Tsung-Yu Aleck Yang ; Taiwan

Taiwania is an open access (available online at <http://taiwania.ntu.edu.tw>) and peer-reviewed journal issued quarterly, with one volume per year. **The Journal** covers all aspects of biodiversity, with a regional focus on East and Southeast Asia. The content is indexed by major abstracting and indexing services, such as SCIE, Biosis Previews, Zoological Record, Scopus, DOAJ and CAB Abstracts. **Author guidelines** are available on the journal website, and shortened version is also on back cover of this issue. Submitted papers must be original and should not be currently submitted nor published elsewhere. Printed copies can be obtained from the **Taiwania Editorial Office, College of Life Science, National Taiwan University, No. 1, Section 4, Roosevelt Road, Taipei 106, Taiwan.**

ISSN: 0372-333X**<https://taiwania.ntu.edu.tw>**

Cover Photo: *Bulbophyllum pulanense* Yudistira, F.H.Kurniawan & Mustaqim, is a new species from Kapuas Hulu Regency, Kalimantan Barat Province of Indonesian Borneo. As a member of the section *Beccariana*, the flowers has a 3-flowered inflorescence, basal pedicel nodes level with bract attachment, lowermost flower's pedicel and ovary more than 2-times length of rachis, entire petals, non-twisted lateral sepals, and tooth or wing absent from the adaxial side of steldia

Acknowledgments: This publication is supported in part by Grants from National Taiwan University and National Science and Technology Council, ROC.